University of Sistan and Baluchestan

Journal of Epigenetics

Online ISSN: 2676 - 4350

Homepage: https://jep.usb.ac.ir

Iranian Genetics Society

Heterosis and combining ability evaluation in different tomato inbred lines for salinity tolerance during the seedling stage

Hamed Kaveh[1*], Safieh vatandoost[2], Hossein Nemati[3], Mohammad Farsi[4]

[1] *Corresponding author*, Assistant professor, Department of Plant Production, University of Torbat Heydarieh. E-mail: h.kaveh@torbath.ac.ir.

[2] Lecturer, Department of Plant Production, University of Torbat Heydarieh. E-mail: s.vatandoost@yahoo.com

[3] Associate professor, Department of horticulture, Ferdowsi University of Mashhad. E-mail: nematish@um.ac.ir

[4] Professor, Department of Biotechnology, Ferdowsi University of Mashhad. E-mail: mofarsi@um.ac.ir

ARTICLE INFO

Article type:
Research Article

Article history:
Received: 01 October 2019
Revised: 01 October 2020
Accepted: 11 October 2020

Keywords:
Cytoplasmic effect,
Reciprocal
cross,
Seedling growth,
Dominance,
Diallel-cross.

A B S T R A C T

Vast numbers of studies on tomato breeding have been performed for resistance to environmental stresses and pests. In genetic investigation and breeding of tomato, it is tried to use tomato lines with the most variance on morphological and agronomical traits to obtain better results, like maximum heterosis on F1 hybrids. In this survey combining ability and heterosis of 7 inbred lines (ME, KaLN3, Fla, CT6, LA3770, R2-05, and DB which was selected from 133 tomato lines of the Ferdowsi University of Mashhad collection, based on variation in morphological, biological and agronomical traits) of tomato and their F1 hybrids were investigated for salinity tolerance during seedling stage using the diallel-cross. Therefore, seed germination percentage and seedling emergence percent and rate under salt stress examined in a complete randomized design with four replications. Results show that the reciprocal effects and SCA were significant for all traits on probability of 1 %. Analysis of Wr-Vr regression also shown that seed germination percentage and seedling emergence percentage and rate under salt stress controlled by over-dominance effects. Best line for resistance to salt stress in juvenility was R205 which has the most GCA for all three traits connected to salt resistance. It may be beneficial to combine this line as a parental line for salinity tolerance increasing of tomato seedlings in breeding programs.

Journal of Epigenetics

Introduction

Tomato (*Solanum lycopersicom*) (2n=2x=24) is one of the most important solanaceous vegetable crops of Peru-Ecuador origin. It is a moderate nutritional crop and is considered as an important source of vitamin A, vitamin C, and minerals. Use of F1 hybrids is one of the means ever-increasing demand for tomato the demand (Peralta et al., 2005; Nuez, 2008). As F1 hybrids in tomato offer several advantages such as earliness, higher productivity, improved quality, resistance to biotic and abiotic stresses and also provide a quick and convenient way of combining desirable characters (Omholt, 2000).

It has been realized that hybrids perform better than self-pollinated cultivars in respect of uniformity of product, yield, quality, resistance to pests and diseases, Adoption to different environmental condition and long storage life

DOI: 10.22111/jep.2020.31828.1017
© Hamed Kaveh
Publisher: University of Sistan and Baluchestan

How to Cite: Kaveh, H., vatandoost, S., Nemati, H., Farsi, M. (2021). Heterosis and Combining Ability Evaluation in Different Tomato Inbred-lines for Salinity Tolerance During the Seedling Stage. Journal of Epigenetics, 2(1), 7-14. DOI: 10.22111/jep.2020.31828.1017

(Titok, 1994). The heterosis in tomato has been exploited for developing hybrids through hand emasculation and hand pollination (Omholt, 2000).

Availability of enough irrigation water with acceptable quality in arid and semi-arid regions of the world, limiting agriculture activities. (Munns, 2002; Al-Busaidi, 2009). While crop productivity reduces by high soil salinity level in these areas and salt susceptible vegetable crops either die or have a yield or growth reduction (Kaveh, 2011). High concentrations of salt (150 mM NaCl and 15 mM CaCl2) in the germination media delays germination and reduce its rate, significantly (Foolad, 1997; Foolad, 1999; Foolad, 2004; Kaveh, 2011). In different developmental stages, tomato regulates different response to salt stress. Usually shoot growth and vigor of younger seedlings become stunted while salinity increases (Foolad, 2004; Li et al., 2011). Zhang et al. reported that the occurrence of a slight saline condition during seedling stage increases final plant height and fresh biomass compare to same saline condition in flowering or fruiting stage (Zhang, 2017).

Genetic transformation used as a tool in tomato breeding for higher salinity tolerance. These studies resulted that the expression of one gene could increase tomato tolerance to salinity conditions (Arrillaga, 1998; Gisbert, 2000; Moghaieb, 2000; Zhang, 2001).

In this study we aimed to find best possible line for salinity tolerance, which can be used in breeding programs.

Materials and methods

Plant material

In this study, seven Lines ME, KaLN3, Fla, CT6, LA3770, R2-05, and DB selected from 133 tomato lines of the Ferdowsi University of Mashhad collection, based on variation in morphological, biological and agronomical traits.

Crossing method

After choosing the parental lines, six plants of each line cultivated in separate rows with a distance of 80 cm between rows and 50 cm between the plants on each row in greenhouse condition. Hand emasculation and hand pollination of different tomato lines were done after flowering (Table1). Pollinated flowers covered with cover sheets to prevent unwanted pollinations.

Table 1. Crossing method of different tomato lines.

	ME	**KALN$_3$**	**Fla**	**CT$_6$**	**LA$_{3770}$**	**R$_{205}$**	**DB**
ME	ME* ME	ME* KALN$_3$	ME * Fla	ME* CT$_6$	ME* LA$_{3770}$	ME* R$_{205}$	ME*7 DB
KAL	KALN$_3$*	KALN$_3$*	KALN$_3$*	KALN$_3$*	KALN$_3$*	KALN$_3$*	KALN$_3$*
N$_3$	ME	KALN$_3$	Fla	CT$_6$	LA$_{3770}$	R$_{205}$	DB
Fla	Fla * ME	Fla* KALN3	Fla * Fla	Fla * CT$_6$	Fla* LA$_{3770}$	Fla * R$_{205}$	Fla * DB
CT$_6$	CT$_6$* ME	CT$_6$* KALN$_3$	CT$_6$ * Fla	CT$_6$* CT$_6$	CT$_6$* LA$_{3770}$	CT$_6$* R$_{205}$	CT$_6$* DB
LA$_{3770}$	LA$_{3770}$* ME	LA$_{3770}$* KALN$_3$	LA$_{3770}$ * Fla	LA$_{3770}$* CT$_6$	LA$_{3770}$* LA$_{3770}$	LA$_{3770}$ * R$_{205}$	LA$_{3770}$ * DB
R$_{205}$	R$_{205}$* ME	R$_{205}$* KALN$_3$	R$_{205}$ * Fla	R$_{205}$* CT$_6$	R$_{205}$* LA$_{3770}$	R$_{205}$* R$_{205}$	R$_{205}$* DB
DB	DB * ME	DB * KALN$_3$	DB * Fla	DB* CT$_6$	DB* LA$_{3770}$	DB* R$_{205}$	DB * DB

Journal of Epigenetics

Seed extraction

Full ripe tomato fruits were collected, and seed extraction was done using fermentation method (Nemati, 2010). Seeds and attached fruit pulp were crushed in a non-metalic container and kept as such for fermentation for 48 hours. During fermentation the seeds get detached from the adhering pulp and settles to the bottom of the container. Then, seeds being washed and dried to 12-14% moisture level.

Germination

Seeds were plated into Petri dishes under aseptic conditions. Each Petri dish contains 20 seeds of one specific cross. Petri dishes were placed in a controlled-environment growth chamber at 25 °C and 80% relative humidity. Irrigation was begun immediately after by adding 20 mL of NaCl (0.5, 2.5, 5, and 10 ds·m−1) solutions daily. To avoid aggregation of salt in dishes, irrigation with distilled water applied every day. When the radical was at least 2 mm long, seeds were considered germinated (Al-Harbi, 2008).

Emergence and Seedling Growth

Tests were carried out in seedling trays with 190 cells (4 ×4 × 6 cm), which were filled with sterilized coco peat and sand 1:1 v/v and placed in a greenhouse at 25 ± 3°C 60% ± 15% relative humidity. Seeds of the tomato Inbred-lines and their cross offspring have sawn one seed per cell. Then, Irrigation was begun immediately after by adding 200 mL of NaCl (0.5, 2.5, 5, and 10 ds·m−1) solutions daily. Irrigation with distilled water repeated every two days to avoid aggregation of salt in the medium. A seedling considered emerged when the hypocotyls hook was visible above the media surface. The emerged seeds number was recorded daily (emergence rate), and after ten days the final emerged seedling percentage determined. The experiment was a factorial, with completely randomized design, with four replications.

Data Analysis

Analysis of Diallel cross for general and specific combining ability and heterosis of each studied characteristics done using Dial 98 (Nemoto et al., 2004).

Results and Discussion

Germination:

ANOVA for germination under saline conditions based on Griffing (1956) revealed that combining ability and cytoplasmic effects are significant (P≤0.01). Progeny of CT6 × LA3770, CT6 × R205, and Fla × R205 crosses with 64,62 and 62, respectively, had the highest percentage of germination. For this trait average of F1 hybrids had a 1.92 fold more than the best parent. CT6 × R205 had the highest amount of heterosis (Table 2). Highest value for SCA has happened in CT6 × LA3770 (Table 3).

Due to the lack of significance in the GCA/SCA ratio in F test, it could be said that additive effects had no significant effects on germination percentage. According to figure 1 and average dominance of this trait (1.201), over dominance effects are more responsible for germination percentage in saline condition.

Table 2. Heterosis value for germination percentage of each cross progeny.

	Me	**KalN3**	**Fla**	**CT6**	**LA3770**	**R205**	**DB**
Me	-	20.5	-2.5	33.75	-8.25	32	-2.75
KaLN3	4	-	-3.5	3.75	-16.75	24	15.25
Fla	-20	-7.5	-	-18.25	0.25	38	0.75
CT6	-3.25	10.75	5.25	-	30.5	52.25	-2.5
LA3770	-8.75	-5.25	-20.25	-7.5	-	-8.75	-14.5
R205	21	24	20	4.75	-5.25	-	21.25
DB	9.25	4.25	-18.25	-1	-25	5.25	-

Table 3. Specific combining ability of germination percentage

	Me	KalN3	Fla	CT6	LA3770	R205	DB	σSij^2
Me	-2.43	2.3	-9.55	1.75	-1.25	4.65	2.1	22.03522381
KaLN3	-2.7	-3.98	-3.75	-6.2	-3.7	2.7	8.65	27.86208095
Fla	-5.9	-2.55	1.87	-8.3	8.95	10.85	1.8	63.03622381
CT6	-4.25	-0.15	5.9	2.82	15.25	3.9	-6.4	67.65689048
LA3770	2.35	1.2	2.5	2.15	2.57	-17.6	-1.65	107.6768905
R205	4.65	3.75	2.05	-3.55	-5.45	9.17	-4.5	93.98853333
DB	5.85	0.45	-2	-0.1	-2.75	-1.45	-10.03	38.77239048
σSij^2	21.0986	7.18291	15.5954	12.8244	10.1269	26.0619	22.3882	

Researchers have reported different inheritance method for the germination percentage. Some researchers have reported that only additive effects play a role in controlling seed germination (Farzaneh,2012). While the results of research by Sekhar et al. (2010) showed that non-additive effects are responsible for inheritance of this trait. Salinity tolerance was mainly controlled by additive effects with the tolerance allele showing partial dominance (Chen, et al; 2008). When High positive SCA was also found between two tolerant cultivars, it is indicating possible different tolerant genes or some minor genes in these cultivars.




Fig.1- Wr-Vr diagram of germination percentage, Wr = aVr + b.

1-ME, 2-KaLN3, 3-Fla, 4-CT6, 5-LA3770, 6- R205, 7-DB.

Seedling emergence

Analysis of variance for seedling emergence showed that maternal effects and SCA were significant (P≤1%) (Table 4).

Table 4. Analysis of variance for seedling emergence based on Griffing (1956)

	df	SS	F	
Block	3	1.71	0.63 $^{n.s}$	(Rep/Error)
GCA[1]	6	710.33	1.17 $^{n.s}$	(GCA/SCA)
SCA[2]	14	608.59	222.22 **	(SCA/Error)
Maternal effects	21	462.29	168.8 **	(Rec/Error)
Error	123	2.74		

n.s: not significant
**: significant at P≤1%.
[1,2]:General and SpecificCombining ability

The total mean of F1 hybrids was higher than best parent value 7.5 units that show heterosis for this trait. Progeny of R205 × CT6 (42.5) and CT6 × KaLN3 (39.5) had the highest value for heterosis for this trait (Table 5). Maximum SCA for this trait in direct crosses went to Me progenies, and for reciprocal crosses, it went to R205 hybrids (Table 6).

Table 5. Heterosis value for emergence percentage of each cross progeny.

	Me	KalN3	Fla	CT6	LA3770	R205	DB
Me	-	-3.75	-6	7.25	-6.75	19.25	16.25
KaLN3	29.25	-	-2.25	12.5	-21.5	-3	18
Fla	-11	-8.25	-	15.25	0.25	26.75	16.25
CT6	1.75	39.5	11.25	-	-11	22.5	-6
LA3770	-6.75	34	18.75	1	-	32.5	-18.5
R205	-2.25	-5	22.25	42.5	-11	-	3
DB	20.25	19	-5.75	2	11	1.5	-

Table 6, Specific combining ability for emergence percentage

	Me	KalN3	Fla	CT6	LA3770	R205	DB	σSij^2
Me	-14.15	12.88	-10.07	7.63	21.54	-10.80	-21.17	254.30
KaLN3	6.65	-2.68	10.61	-13.64	1.54	-8.51	-2.87	92.24
Fla	-11.50	-13.35	0.75	8.54	-3.69	-13.56	8.17	91.92
CT6	-4.60	11.80	2.15	-2.90	-14.51	13.46	-1.49	119.72
LA3770	-4.20	3.70	10.05	-10.55	-12.61	-1.42	-3.46	140.09
R205	-1.65	-19.25	12.35	14.25	4.15	13.35	20.82	190.38
DB	15.30	10.45	0.30	-13.05	-3.15	-9.85	18.25	209.37
σSij^2	103.94	143.55	94.87	109.49	67.86	165.97	151.00	

The average degree of dominance which is equal to 1.44 and b value of figure 2 (-65.97) showed that overdominance controlling this trait in most of gene loci.

For emergence percentage, there was no specific research on Tomato, but Zengin (2016) described genetic structure of the 30 F1 hybrid tomato combinations and Two Male Testers. It has been suggested that both additive and non-additive effects control this trait.




Fig. 2. Wr-Vr diagram of emergence percentage, Wr = aVr + b.
1-ME, 2-KaLN3, 3-Fla, 4-CT6, 5-LA3770, 6- R205, 7-DB.

Seedling growth

Analysis of variance for this trait (Table 7) showed that general and specific combining ability and maternal effects were significant.

The average of heterosis for this trait was equal to -3.3. Although there were high positive heterosis values in this trait for some of the inbred-lines like R205 and LA3770 negative heterosis was more favorable. As much as heterosis value became more negative, the time that seedling reached to six leaves-stage was reduced. This means that progeny of LA3770 and CT6 had the fastest growth rate during seedling stage (Table 8).

Table 7. Analysis of variance for seedling emergence rate based on Griffing (1956)

	df	SS	F	
Block	3	105.33	0.63 $^{n.s}$	(Rep/Error)
GCA	6	4220.72	2.04 *	(GCA/SCA)
SCA	14	2068	12.46 **	(SCA/Error)
Maternal effects	21	2413.33	14.54 **	(Rec/Error)
Error	123	165.97		

n.s: not significant
*: significant at P≤5%.
**: significant at P≤1%.

Table 8, heterosis value for emergence rate of each cross progeny

	Me	KalN3	Fla	CT6	LA3770	R205	DB
Me	-	-15.85	-30.1	17.25	-24.55	-24.8	-0.5
KaLN3	-28.85	-	21.55	-25.6	18.1	65.85	-6.85
Fla	-43.1	-77.95	-	-37.35	20.35	51.6	3.9
CT6	-58.25	-51.6	43.15	-	-65.3	46.95	-53.25
LA3770	7.95	-20.4	-58.65	-31.3	-	18.65	40.95
R205	21.7	53.35	-29.4	51.95	8.15	-	-3.3
DB	-18	9.65	2.4	32.75	19.45	10.7	-

In this trait, lower values of specific combining ability means that the F1 hybrid was more resistant to saline conditions, so CT6 × LA3770 and R205 × DB hybrids were the best (Table 9).

Table 9, Specific combining ability for emergence rate

	Me	KalN3	Fla	CT6	LA3770	R205	DB	σSij^2
Me	-24.83	-1.40	-8.15	7.65	13.45	-13.60	2.05	170.90
KaLN3	3.80	8.37	-13.60	-24.30	6.75	33.70	-1.15	336.67
Fla	1.45	-16.25	-11.88	24.70	-3.75	-7.30	8.10	185.90
CT6	-21.15	-18.10	34.05	-4.43	-33.20	30.75	-5.60	559.18
LA3770	18.10	3.65	-10.70	-12.05	5.27	-11.70	28.45	385.48
R205	5.85	21.40	-15.20	10.45	-5.30	16.32	-31.85	616.52
DB	-8.05	5.50	6.65	6.80	6.30	-17.20	11.17	338.88
σSij^2	236.49	194.62	322.72	375.30	115.37	230.77	104.66	




Figure 3- Wr-Vr diagram of emergence rate, Wr = aVr + b.
1-ME, 2-KaLN3, 3-Fla, 4-CT6, 5-LA3770, 6- R205, 7-DB.

Analysis of figure 3 showed that b value was equal to -452.866. Considering b value and average dominance degree (2.79) we can conclude that over dominance may have a big role in controlling this trait in saline conditions.

Conclusion

For germination percentage and seedling emergence and growth in studied tomato lines we had concluded that R205 ،CT6 ،LA3770 و KaLN3 and their progeny had better performance in saline condition, and the best line for tolerance to salinity during seedling stage was R205 which had a high amount of GCA in total, and it's progeny, or one of them performed better than others. It may be beneficial to combine this line as a parental line for salinity tolerance increasing of tomato seedlings in breeding programs.

References

Al-Busaidi, A., Al-Rawahy, S., Ahmed, M., 2009. Response of Different Tomato Cultivars to Diluted Seawater Salinity. Asian Journal of Crop Science 1, 77-86.

Al-Harbi, A.R., Wahb-Allah, M. A., Abu-Muriefah, S. S., 2008. 'Salinity and Nitrogen Level Affects Germination, Emergence, and Seedling Growth of Tomato. International Journal of Vegetable Science 14, 380-392.

Arrillaga, I., Gil Mascarell, R., Gisbert, C., Sales, E., Montesinos, C., Serrano, R., and Moreno, V., 1998. Expression of the yeast HAL2 gene in tomato increases the in vitro salt tolerance of transgenic progenies. plant science 136, 219-226.

Chen, Z., Shabala, S., Mendham, N., Newman, I., Zhang, G. and Zhou, M. (2008), Combining Ability of Salinity Tolerance on the Basis of NaCl-Induced K+ Flux from Roots of Barley . Crop Science, 48: 1382-1388. doi:10.2135/cropsci2007.10.0557.

Farzaneh, Akram. (2012). Estimation of the ability to combine and heterozygous for valuable agricultural traits in nine tomato lines using the dial method. Master Thesis, Ferdowsi University of Mashhad.

Foolad, M.R., 1997. Genetic basis of physiological traits related to salt tolerance in tomato, Lycopersicon esculentum Mill. Plant Breeding 116, 53-58.

Foolad, M.R., 1999. Comparison of salt tolerance during seed germination and vegetative growth in tomato by QTL mapping. Genome 42, 727-734.

Foolad, M.R., 2004. Recent Advances in Genetics of Salt Tolerance in Tomato. Plant Cell, Tissue and Organ Culture 76, 101-119.

Gisbert, C., Rus, A.M., Bolarín, M.C., Coronado, J.M., Arrillaga, I., Montesinos, C., Caro, M., Serrano, R., and Moreno, V. , 2000. The yeast HAL1 gene improves salt tolerance of transgenic tomato. plant physiology 123.

Kaveh, H., Nemati,H., Farsi, M., and Vatandoost Jartoodeh, S., 2011. How Salinity Affect Germination and Emergence of Tomato Lines. J. BIOL. ENVIRON. SCI. 5, 159-163.

Li, J., Liu, L., Bai, Y., Zhang, P., Finkers, R., Du, Y., Visser, R., van Heusden, A., 2011. Seedling salt tolerance in tomato. Euphytica 178, 403-414.

Moghaieb, R.E.A., Tanaka, N., Saneoka, H., Hussein, H.A., Yousef, S.S., Ewada, M.A.F., Aly, M.A.M., and Fujita, K., 2000. Expression of betaine aldehyde dehydrogenase gene in transgenic tomato hairy roots leads to the accumulation of glycine betaine and contributes to the maintenance of the osmotic potential under salt stress. soil science plant nutrition 46, 873-883.

Munns, R., 2002. Comparative physiology of salt and water stress. Plant, Cell & Environment 25, 239-250.

Nemati, H., Nazdar,T., Azizi,M., and Arouiee,H., 2010. The Effect of Seed Extraction Methods on Seed Quality of Two Cultivar's Tomato (Solanum lycopersicum L.). Pakistan Journal of Biological Sciences 13, 814-820.

Nemoto, K., Ukai, Y., Tang, D.Q., Kasai, Y., Morita, M., 2004. Inheritance of early elongation ability in floating rice revealed by diallel and QTL analyses. TAG Theoretical and Applied Genetics 109, 42-47.

Nuez, F., and Prohens.,J. 2008. handbook of plant breeding.vegetables II. (Ed), 2008. Hand book of plant breeding, Vegetables II. Springer, New york.

Omholt, S.W., Plahte, E., Oyehaug, L., and KeFang, X., 2000. Gene regulatory networks generating the phenomena of additivity, dominance and epistasis. Genetics 155, 969-980.

Peralta, I.E., Knapp, S., Spooner, D.M., 2005. New Species of Wild Tomatoes (Solanum Section Lycopersicon: Solanaceae) from Northern Peru. Systematic Botany 30, 424-434.

Sekhar, L., Prakash, B.G., Salimath, P.M., Channayya, M., Hiremath, P., Sridevi, O. and Patil, A.A., 2010. Implication of heterosis and combining ability among productive Single cross Hybrid in tomato. Electronic Jornal of Plant Breeding, 1(4): 706-711.

Titok, V.V., Lemesh, V.A., Rusinova, O.V., and Podlisskikh, V.L., 1994. Leaf area, chlorophyll content and biomass of tomato plants and their heterotic hybrids under in vitro culture. . Photosynthetica 30, 255-260.

Zengin, S., KABAŞ, A., OĞUZ, A., EREN, A., POLAT, E., 2016. Determining of general combining ability for yield, quality and some other traits of tomato (Solanum lycopersicum L.) inbred lines. Akdeniz Üniversitesi Ziraat Fakültesi Dergisi , 28 (1) , 0-0 . Retrieved from https://dergipark.org.tr/en/pub/akdenizfderg/issue/25317/267420

Zhang, H.X., Blumwald, E. , 2001. Transgenic salt-tolerance tomato plants accumulate salt in foliage but not in fruit. Nature Biotechnology 19.

Journal of Epigenetics